Evidence-Based Nursing



Contents

Purpose and procedure

65

EBN notebook

67 Economic evaluation of healthcare technologies using primary research

Treatment

- **72** Review: venepuncture is less painful than heel lance for blood sampling in neonates
- 73 Review: parenting interventions can reduce unintentional injuries in children and increase use of safety practices
- 74 Review: cranberry products may prevent urinary tract infection in women with recurrent infections
- **75** Review: use of pedometers increases physical activity in adults
- **76** Telling smokers their "lung age" promoted successful smoking cessation
- 77 Review: behavioural interventions promote tobacco cessation in users of smokeless tobacco
- 78 Review: antibiotics are only slightly more effective than placebo for clinically diagnosed acute rhinosinusitis in adults
- 79 Therapeutic Interactive Voice Response enhanced CBT gains in chronic pain
- **80** Review: evidence is lacking on the effectiveness of dietary advice in illness-related malnutrition
- 81 Spontaneous awakening trials plus spontaneous breathing trials improved weaning from mechanical ventilation
- **82** Review: prone positioning improved oxygenation on ventilation more than supine positioning in ARDS
- 83 Backfill assisted voiding was better than spontaneous voiding for successful bladder emptying after vaginal surgery
- **84** Review: organised stroke unit care reduces mortality more than other forms of care for inpatients
- **85** Review: poor-quality trials suggest that some interventions for family caregivers of patients with stroke improve carer outcomes

July 2008 Vol 11 No 3

86 Telehome monitoring reduced readmissions and improved quality of life in heart failure or angina



Honey-impregnated dressings and usual care did not differ for healing venous leg ulcers

Review: evidence for use of systemic antibiotics and topical antiseptics for venous leg ulcers is insufficient

Diagnosis

89 Chlamydia Rapid Test was moderately accurate for diagnosing Chlamydia infection in women

Causation

90 Review: use of cannabis is associated with increased risk of psychotic outcomes later in life

Clinical prediction guide

91 QRISK underestimated risk of cardiovascular disease in general practice patients; Framingham score overestimated risk

Qualitative

- 92 Adolescents learned self-management of arthritis by acquiring knowledge and skills and experiencing understanding from social support
- 93 3 themes described what involvement in treatment decision making meant to patients with diabetess
- 94 Women with spinal cord injuries underwent a process of discomfort, moving towards comfort, and comfort in dealing with their changed bodies
- 95 A potential barrier to discussing deactivation of implantable cardioverter defibrillators was patients' lack of knowledge
- 96 3 patterns described the meaning of memory loss within everyday life for people with early Alzheimer disease

Glossary

Inside back cover





