
# **Evidence-Based Nursing**



# **Contents**

#### **Editorial**

97 Patient and public involvement in the implementation of evidence into practice

#### Research made simple

**98** Understanding triangulation in research

#### Midwifery

99 Cohort study finds newborn respiratory complications less common when mothers of babies with fetal lung immaturity at 34–37 weeks' gestation given antenatal steroids

#### **Adult nursing**

101 Exercise improves fatigue during and after breast and prostate cancer treatment, with benefits seen for aerobic exercise

#### Child health

- Contact with dogs during the first year of life is associated with decreased risk of respiratory illness
- 104 For vulnerable families, continued postnatal care, provided by family midwives, is associated with improved maternal care and parent—child relationship
- 105 Vitamin D concentration in newborn cord blood is correlated with maternal intake of supplemental vitamin D, and lower levels are associated with increased risk of the infant developing eczema

#### **Health promotion**

- 107 Sunbed use increases risk of melanoma; risk increases with greater number of sessions and first use at younger age
- Mobile phone-based smoking cessation interventions increase long-term quit rates compared with control programmes, but effects of the interventions are heterogeneous

#### **Primary health care**

- People with diabetes and self-reported severe hypoglycaemia have increased mortality risk over 5 years
- 111 General health checks may not reduce morbidity or mortality but do increase the number of new diagnoses

### October 2013 Vol 16 No 4

Compared to standard physician care for diabetes, nurse specialists provide similar care in terms of quality of life

113

Updated systematic review suggests that cranberry juice is not effective at preventing urinary tract infection

# Pain management

Movement pain, resting pain and depression prior to total knee replacement predict postoperative pain

## Nursing issues

- 117 Scientific nursing journals over 25 years: most studies continue to focus on adults and psychological variables, with a decline in theory-testing-based studies and an increase in qualitative studies
- 118 US Medicare data show incidence of hospital-acquired pressure ulcers is 4.5%, and they are associated with longer hospital stay and higher risk of death
- 119 Hospital nursing units in the USA: government ownership, Magnet designation, unit population age group and higher skill mix are associated with lower nursing turnover
- 120 Cochrane reviews allow firm conclusions to be made about the effectiveness of various local and systemic treatments for wounds and ulcers

#### Care of older people

- 122 Increased city-level influenza mortality and statelevel influenza severity is associated with greater functional decline among nursing home residents
- 123 Pain assessment and management in older hospitalised patients: observation shows common themes relating to nurse—patient communication, pain management strategies, organisational aspects of care and the complex nature of pain
- 124 Exposure to challenging behaviour from nursing home residents is associated with reduced general health and work ability, and increased burnout reported by nurses

#### **PostScript**

**126** Letter of response to the commentary written by Dr Howell





Receive regular table of contents by email. Register using this QR code.