Available online in April at www.evidencebasednursing.com

## **Evidence-Based Nursing**

## Contents

Purpose and procedure34	Integrated care pathway based rehabilitation for acute stroke did not reduce length of hospital stay58
EBN notebook	Intensive geriatric rehabilitation reduced hospital stay and time to independent living in hip fracture patients with mild to moderate dementia
Clinically useful measures of the effects of treatment36	
Sigma Theta Tau contest	Assessment
Evidence-based practice contest: innovations in clinical excellence	A clear urine specimen on visual inspection cannot totally exclude a diagnosis of urinary tract infection
Treatment	Causation
Routine primary immunisation using a longer needle resulted in fewer local reactions in infants	Children who were exposed to antenatal corticosteroids and had birth weights $\leq 1500$ g did not have adverse growth, cognitive, or lung function outcomes at age 14 years56
Few young children had mild reactions to the measles, mumps, and rubella vaccine and older children had nearly no vaccine related reactions	Quality improvement
Review: recall or reminder systems improve the rate of childhood and adult immunisation	A nurse led clinic and computer decision support system for anticoagulation decisions was at least as effective as a hospital clinic
A clinical pathway reduced use of nebulised $\beta$ agonists and length of hospital stay in children with asthma exacerbations	Clinical prediction guide
Review: brief primary care interventions are moderately effective for increasing physical activity	An antenatal index predicted postpartum depression with high sensitivity and moderate specificity58
Contamination of urine specimens from women with acute dysuria did not differ with collection technique46	Qualitative
Review: NSAIDs, paracetamol, and analgesic/ anti-inflammatory oral rinse reduce sore throat symptoms 47	Late adolescent girls used 4 stories to describe their smoking patterns59
Docusate sodium with irrigation was better than triethanolamine polypeptide with irrigation for dissolving	Teenage mothers with oppressive pasts created new family caregiving traditions
earwax	Individuals with HIV/AIDS ascribed several different meanings to their use of complementary therapy
Early feeding after major non-laparoscopic gynaecological surgery did not increase gastrointestinal symptoms49	Inpatients had mostly negative experiences of seclusion during short term treatment in a mental health facility65
Adjunctive nurse telehealth care reduced depressive symptoms and improved functioning50	Properly planned de-institutionalisation for mental illness
A shared care system of hospital follow up reduced pain and use of healthcare resources and increased satisfaction in	maintained most in community living with enhanced quality of life
patients with rheumatoid arthritis	Elderly women described 5 phases of living with change after myocardial infarction
A cognitive behavioural family intervention reduced psychiatric morbidity in caregivers of patients with	Glossary