

Evidence-Based Nursing

Contents

Purpose and procedure	98	Assessment (screening or diagnosis)	
Implementation forum		The SCOFF questionnaire was less sensitive but more specific than the ESP for detecting eating disorders	118
Building a foundation for evidence-based practice: experiences in a tertiary hospital	100	Prognosis	
Resource review	104	Needing help to use a toilet was associated with faecal incontinence at 3 months in new onset faecal incontinence after stroke	119
Treatment		Causation	
Providing free smoke alarms did not reduce fire related injuries in a deprived multiethnic urban population	105	Eating alone, parents' marital status, and use of radio and girls' magazines were risk factors for eating disorders	120
Review: self management education improves outcomes in children and adolescents with asthma	106	Intentional weight loss was associated with lower mortality, whereas unintentional weight loss was associated with increased mortality	121
Review: regular medical review is not better than written self management plans for optimising asthma control	107	Quality improvement	
Review: day hospital is as effective as inpatient care but has lower costs for patients with acute psychiatric disorders	108	Preoperative assessments by trained nurses were equal in quality to assessments by preregistration house officers	122
Treatment of hypertension reduces risk of complications in type 2 diabetes	109	The benefits of nurse led secondary prevention clinics for coronary heart disease continued after 4 years	123
An intensive, target driven intervention reduced cardiovascular and microvascular events in patients with type 2 diabetes and microalbuminuria	110	Qualitative	
A COPD self management programme reduced hospital use and improved health status	111	Individuals taking combination therapies for HIV or AIDS faced barriers to remaining in, or returning to work	124
Review: several interventions prevent ventilator associated pneumonia in critically ill patients	112	Living with untreated localised prostate cancer was seen as living under a dark shadow	125
Oral vitamin D ₃ supplementation reduced fractures in community dwelling elderly people	113	Women living with ovarian cancer described changes in day to day living, major challenges, and sources of support	126
A multifactorial intervention after a fall did not prevent falls in elderly patients with cognitive impairment and dementia	114	Women with ovarian cancer described waiting for recurrence, facing the diagnosis of recurrence, managing treatment related concerns, and attempting to regain control	127
A structured education programme increased hip protector use and may reduce hip fractures in nursing homes	115	Parents of adult children with schizophrenia dealt with societal objections in order to provide the necessary care	128
Resistance exercises or vitamin D did not improve physical health or reduce falls in frail older people	116	Glossary	Inside back cover
An exercise and incontinence intervention did not reduce the incidence or cost of acute conditions in nursing home residents	117	Other articles noted (25 Apr 03 to 18 Jul 03)	e1

Search all issues of Evidence-Based Nursing published since 1998 and over 300 medical journals through Highwire press at www.evidencebasednursing.com. EBN 77944.